

DIRAC OPERATORS ON QUANTUM TWO SPHERES

Kazutoshi Ohta* and Hisao Suzuki†

*Department of Physics,
Osaka University
Toyonaka, Osaka 560, Japan*

Abstract

We investigate the spin 1/2 fermions on quantum two spheres. It is shown that the wave functions of fermions and a Dirac Operator on quantum two spheres can be constructed in a manifestly covariant way under the quantum group $SU(2)_q$. The concept of total angular momentum and chirality can be expressed by using q -analog of Pauli-matrices and appropriate commutation relations.

*e-mail address: kohta@oskth.kek.jp

†e-mail address: suzuki@fuji2.wani.osaka-u.ac.jp

The structure of quantum groups [1][2][3] often appears as hidden symmetry in physics. One of interests in the quantum groups is that we can describe quantum manifolds as a non-commutative geometry. [4][5][6][7] The quantum manifolds have been studied as possible deformation of manifolds along with the geometrical understanding of the quantum groups. And that deformation can be used to generalize the ordinary notion of space(time) symmetry.

Recently, it was suggested that compact quantum manifolds can be used as an regularization of field theories. It was shown[8] for the scalar field on the quantum 2, 3 and 4 spheres[5][9] that the number of wave functions can be finite when q is a root of unity ($q^l = 1$). The parameter l is determined by the ratio of the scale of the universe and that of the ultraviolet cut-off. Another attractive feature of the quantum spheres is the scale dependence. The effects of the deformed manifold are invisible at sufficient low energy scale. In other words, the quantum manifolds might (effectively) describe the strong quantum fluctuations of the manifolds in microscopic level, which can be smoothed out in our experimental scale.

Although the scalar fields on quantum spheres are constructed consistently, the consistency of the higher spin fields has not been clear. The construction of the higher spin fields is required for the full realization of the quantum field theories on quantum spheres.

In this letter, we deal with the quantum S^2 and construct Dirac operators. The advantage of treating the quantum two sphere is that this manifold is not only the simplest compact quantum manifold in even dimensions where we can discuss the chiral fermions, but we can use well known properties of representation theories of quantum $SU(2)_q$.

The algebra $SU(2)_q[1][2]$ is generated by elements $X^\pm$ and $K = q^{H_0}$ satisfying the following relations

$$KX^\pm K^{-1} = q^\pm X^\pm, \quad [X^+, X^-] = \frac{K - K^{-1}}{q^{\frac{1}{2}} - q^{-\frac{1}{2}}}. \quad (1)$$

A structure of a Hopf algebra is given by

$$\Delta(X^\pm) = X^\pm \otimes K^{-\frac{1}{2}} + K^{\frac{1}{2}} \otimes X^\pm, \quad \Delta(K) = K \otimes K. \quad (2)$$

Our convention of this structure describe the one of $SU(2)_{q^{-1}}$ in the convention of ref.[10].

The generators of $SU(2)_q$ act on the spin j representation in the following way,

$$X^\pm |j, m\rangle = ([j \mp m][j \pm m + 1])^{\frac{1}{2}} |j, m \pm 1\rangle \quad (3)$$

$$K |j, m\rangle = q^m |j, m\rangle, \quad (4)$$

where $-j \leq m \leq j$, $2m \equiv 2j \pmod{2}$ and,

$$[n] \equiv \frac{q^{\frac{n}{2}} - q^{-\frac{n}{2}}}{q^{\frac{1}{2}} - q^{-\frac{1}{2}}}. \quad (5)$$

We introduce coordinates of quantum sphere[5] as an spin 1 representation of $SU(2)_q$ which will be denoted by e_m , $m = -1, 0, 1$. The commutation relations among coordinates can be defined by requiring that total spin 1 states cannot be constructed out of the product of the coordinates, which implies

$$\sum_{m'} \begin{bmatrix} 1 & 1 & 1 \\ m' & m - m' & m \end{bmatrix}_q e_{m'} e_{m-m'} = 0, \quad (6)$$

where $\begin{bmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{bmatrix}_q$ represents the q analog of Clebsh-Gordan Coefficient[10]. For later convenience, we list the following explicit form:

$$\begin{bmatrix} 1 & j & j \\ 0 & m & m \end{bmatrix}_q = \{q^{(j-m+1)/2}[j+m] - q^{-(j+m+1)/2}[j-m]\} / \sqrt{[2j][2j+2]}, \quad (7)$$

$$\begin{bmatrix} 1 & j & j \\ 1 & m-1 & m \end{bmatrix}_q = -q^{-m/2} \left(\frac{[2][j+m][j-m+1]}{[2j][2j+2]} \right)^{1/2}, \quad (8)$$

$$\begin{bmatrix} 1 & j & j \\ -1 & m+1 & m \end{bmatrix}_q = q^{-m/2} \left(\frac{[2][j-m][j+m+1]}{[2j][2j+2]} \right)^{1/2}. \quad (9)$$

$$(10)$$

The equation of the sphere can be expressed as

$$e \cdot e = r^2, \quad (11)$$

where $a \cdot b$ for spin j representations a_m, b_m is defined by

$$a \cdot b \equiv -\sum_m (-1)^{j-m} q^{m/2} a_{-m} b_m$$

$$= -[2j+1]^{1/2} \sum_m \begin{bmatrix} j & j & 0 \\ -m & m & 0 \end{bmatrix}_q a_{-m} b_m, \quad (12)$$

The highest weight state of integer spin j can be written as $|j, j\rangle = a_j e_1^j$, where the coefficient a_j , which we do not need to specify in this letter, can be determined from the haar measure[3][11][8]. The states $|j, m\rangle$ can be obtained by using a relation (3). For example, the state $|j, j-1\rangle$ is given by $a_j q^{-(j-1)/2} (\frac{[2j]}{[2]})^{1/2} e_1^{j-1} e_0$.

We next introduce the derivatives with respect to the coordinates satisfying

$$\partial_{m_1} e_{m_2} = q^{-1} \sum (R^{1,1})_{m_1, m_2}^{m'_1, m'_2} e_{m'_2} \partial_{m'_1} + (-1)^{m_2} q^{m_2/2} \delta_{m_1+m_2, 0}, \quad (13)$$

where $(R^{1,1})_{m_1, m_2}^{m'_1, m'_2}$ is so-called the universal R-matrix[10] between the spin 1 states. The element can be written in terms of the $3j$ -symbol as follows:

$$\begin{aligned} (R^{j_1, j_2})_{m_1, m_2}^{m'_1, m'_2} &= \sum_j (-1)^{j_1+j_2-j} q^{-\{j_1(j_1+1)+j_2(j_2+1)-j(j+1)\}/2} \\ &\times \begin{bmatrix} j_1 & j_2 & j \\ m_1 & m_2 & m \end{bmatrix}_q \begin{bmatrix} j_2 & j_1 & j \\ m'_2 & m'_1 & m \end{bmatrix}_q, \end{aligned} \quad (14)$$

in which $m_1 + m_2 = m'_1 + m'_2 = m$. Using these derivatives, we can define the operator of orbital angular momentum J_m as

$$J_m = \left(\frac{[4]}{[2]} \right)^{1/2} \sum_{m_1} \begin{bmatrix} 1 & 1 & 1 \\ m_1 & m-m_1 & m \end{bmatrix}_q e_{m_1} \Lambda \partial_{m-m_1}, \quad (15)$$

where Λ is the dilatation operator[12][13][14] whose action to e_m and ∂_m is given by

$$\Lambda e_m = q e_m \Lambda, \quad \Lambda \partial_m = q^{-1} \partial_m \Lambda. \quad (16)$$

The introduction of this operator is required to ensure an relation

$$[J_m, r^2] = 0. \quad (17)$$

Since each operator e_m and ∂_m changes the spin of the state by ± 1 , the operator J_m does not change the spin of the states. (The total change of spin by ± 2 is forbidden.)

Therefore, the operators J_m can be called as quantum rotation operators. The action of this operator on the states $|j, m\rangle$ can be specified as

$$J_m|j, m'\rangle = \frac{[2j]^{1/2}[2j+2]^{1/2}}{[2]} \begin{bmatrix} 1 & j & j \\ m & m' & m+m' \end{bmatrix}_q |j, m+m'\rangle, \quad (18)$$

where we have obtained the coefficient by acting J_0 to the highest weight state because the m dependence should be included only in the $3j$ -symbols. We can define a second order Casimir operator by using J_m as

$$C = J \cdot J, \quad (19)$$

where the product is defined in (12). By using (18) and relations

$$\begin{bmatrix} j_1 & j_2 & j \\ m_1 & m_2 & m \end{bmatrix}_q = (-1)^{j_2-j-m_1} q^{-m_1/2} \left(\frac{[2j+1]}{[2j_2+1]} \right)^{1/2} \begin{bmatrix} j_1 & j & j_2 \\ -m_1 & m & m_2 \end{bmatrix}_q, \quad (20)$$

$$\sum_{m_1, m_2} \begin{bmatrix} j_1 & j_2 & j \\ m_1 & m_2 & m \end{bmatrix}_q \begin{bmatrix} j_1 & j_2 & j' \\ m_1 & m_2 & m' \end{bmatrix}_q = \delta_{j,j'} \delta_{m,m'}, \quad (21)$$

for $|j_1 - j_2| \leq j \leq j_1 + j_2$, we find the value of the Casimir operator as

$$C|j, m\rangle = \frac{[2j][2j+2]}{[2]^2} |j, m\rangle. \quad (22)$$

Using this Casimir operator, we can construct an equation of the scalar field of mass m on this Euclidean quantum space as

$$\left(\frac{C}{r^2} + m^2 \right) \phi(e) = 0. \quad (23)$$

Our next question is whether chiral fermions can exist on this quantum two sphere.

To begin with, we operatorially introduce the spin $1/2$ spinors as

$$\begin{aligned} |1/2, 1/2\rangle &= c_{1/2} |0\rangle, \\ |1/2, -1/2\rangle &= c_{-1/2} |0\rangle. \end{aligned} \quad (24)$$

It is not necessary to introduce commutation rule among the operators c 's since we only consider one particle states with respect to c 's.. We next define the conjugate of the fields c_m preserving the $SU(2)_q$ invariant:

$$b_{m_1} c_{m_2} |0\rangle = [2]^{1/2} \begin{bmatrix} \frac{1}{2} & \frac{1}{2} & 0 \\ m_1 & m_2 & 0 \end{bmatrix}_q |0\rangle. \quad (25)$$

Explicitly, we obtain

$$b_{-1/2} c_{1/2} |0\rangle = q^{\frac{1}{4}} |0\rangle, \quad b_{1/2} c_{-1/2} |0\rangle = -q^{\frac{1}{4}} |0\rangle, \quad (26)$$

$$b_{1/2} c_{1/2} |0\rangle = b_{-1/2} c_{-1/2} |0\rangle = 0. \quad (27)$$

From these operators, we can introduce q-analog of Pauli matrices as

$$\sigma_m = [2]^{1/2} \sum_{m_1} \begin{bmatrix} \frac{1}{2} & \frac{1}{2} & 1 \\ m_1 & m - m_1 & m \end{bmatrix}_q c_{m_1} b_{m-m_1}, \quad (28)$$

which implies

$$\begin{aligned} \sigma_1 &= [2]^{1/2} c_{1/2} b_{1/2}, \\ \sigma_0 &= q^{-1/4} c_{-1/2} b_{1/2} + q^{1/4} c_{1/2} b_{-1/2}, \\ \sigma_{-1} &= [2]^{1/2} c_{-1/2} b_{-1/2}. \end{aligned} \quad (29)$$

We also define a singlet operator as

$$\begin{aligned} O &= c \cdot b \\ &= -q^{1/4} c_{-1/2} b_{1/2} + q^{-1/4} c_{1/2} b_{-1/2}. \end{aligned} \quad (30)$$

The action of this operator on $c_{\pm}|0\rangle$ is just an identity operator. The q -deformed Pauli-matrices form an algebra:

$$\sigma_{m_1} \sigma_{m_2} = \frac{[4]^{1/2}}{[2]^{1/2}} \begin{bmatrix} 1 & 1 & 1 \\ m_1 & m_2 & m_1 + m_2 \end{bmatrix}_q \sigma_{m_1+m_2} + (-1)^{m_1} q^{-m_1/2} \delta_{m_1+m_2,0} O, \quad (31)$$

which is the q-analog of the well known relation $\sigma^i \sigma^j = i \varepsilon_{ijk} \sigma^k + \delta^{ij}$.

Since the general states can be expressed as products of the coordinates and c_m , we need to specify the commutation relations between e_m and $c_{m'}$. A consistent choice is to use the universal R-matrix[10] with respect to spin 1/2 and spin 1:

$$c_{m_1} e_{m_2} = (R^{1/2,1})_{m_1, m_2}^{m'_1, m'_2} e_{m'_2} c_{m'_1}. \quad (32)$$

We can find that c_m commutes with the radius r^2 . As for the commutation relation between b_{m_1} and e_{m_2} , we prepare two type of conjugate of c_m . One is defined to commute with e_m by R^{-1} matrix which we call b_m , the other is to commute with e_m by R-matrix, which will be b'_m . Correspondingly, the Pauli-matrices and operator (30) constructed by c_m and b_m will be denoted by σ_m and O and the operators constructed by c and b'_m will be called σ'_m and O' .

It seems instructive to write down the commutation relations explicitly:

$$\begin{aligned} c_{1/2} e_1 &= q^{1/2} e_1 c_{1/2}, & c_{-1/2} e_{-1} &= q^{1/2} e_{-1} c_{-1/2}, \\ c_{1/2} e_0 &= e_0 c_{1/2} + [2]^{1/2} q^{-1/4} (q^{1/2} - q^{-1/2}) e_1 c_{-1/2}, \\ c_{-1/2} e_1 &= q^{1/2} e_1 c_{-1/2}, & c_{-1/2} e_0 &= e_0 c_{-1/2}, \\ c_{1/2} e_{-1} &= q^{-1/2} e_{-1} c_{1/2} + [2]^{1/2} q^{-1/4} (q^{1/2} - q^{-1/2}) e_0 c_{-1/2}, \\ b_{1/2} e_1 &= q^{-1/2} e_1 b_{1/2}, & b_{-1/2} e_{-1} &= q^{-1/2} e_{-1} b_{-1/2}, \\ b_{-1/2} e_0 &= e_0 b_{-1/2} - [2]^{1/2} q^{1/4} (q^{1/2} - q^{-1/2}) e_{-1} b_{1/2}, \\ b_{1/2} e_{-1} &= q^{1/2} e_{-1} b_{1/2}, & b_{1/2} e_0 &= e_0 b_{1/2}, \\ b_{-1/2} e_1 &= q^{1/2} e_1 b_{-1/2} - [2]^{1/2} q^{1/4} (q^{1/2} - q^{-1/2}) e_0 b_{1/2}, \end{aligned} \quad (33)$$

The states constructed by the product of spin j representation of the coordinate space and spin 1/2 representation split into the total spin $j + 1/2$ and spin $j - 1/2$. The highest weight states can be written as

$$|j, j + 1/2, j + 1/2\rangle = |j, j\rangle c_{1/2} |0\rangle, \quad (35)$$

$$\begin{aligned} |j, j - 1/2, j - 1/2\rangle &= \\ \frac{1}{[2j + 1]^{1/2}} \{ & q^{j/2} |j, j - 1\rangle c_{1/2, 1/2} - q^{-1/4} [2j]^{1/2} |j, j\rangle c_{-1/2} \} |0\rangle. \end{aligned} \quad (36)$$

The other descendants can be derived by using a relation (3).

We are going to consider the action of σ_m on these states. Since σ_m does not change bosonic spin and preserve $SU(2)_q$ invariance, we have

$$\begin{aligned} \sigma_m |j, j + 1/2, m' \rangle = & b_{++} \begin{bmatrix} 1 & j + 1/2 & j + 1/2 \\ m & m' & m + m' \end{bmatrix}_q |j, j + 1/2, m + m' \rangle \\ & + b_{+-} \begin{bmatrix} 1 & j + 1/2 & j - 1/2 \\ m & m' & m + m' \end{bmatrix}_q |j, j - 1/2, m + m' \rangle. \end{aligned} \quad (37)$$

The coefficients b_{++} and b_{+-} can be determined by considering $\sigma_0 |j, j + 1/2, j + 1/2 \rangle$ and $\sigma_{-1} |j, j + 1/2, j + 1/2 \rangle$, and we get

$$b_{++} = \left(\frac{[2j + 3]}{[2j + 1]} \right)^{1/2} q^j, \quad (38)$$

$$b_{+-} = - \left(\frac{[2j + 2][2]}{[2j + 1]} \right)^{1/2} q^{-1/2}. \quad (39)$$

Quite similarly, we have

$$\begin{aligned} \sigma_m |j, j - 1/2, m' \rangle = & b_{-+} \begin{bmatrix} 1 & j - 1/2 & j + 1/2 \\ m & m' & m + m' \end{bmatrix}_q |j, j + 1/2, m + m' \rangle \\ & + b_{--} \begin{bmatrix} 1 & j - 1/2 & j - 1/2 \\ m & m' & m + m' \end{bmatrix}_q |j, j - 1/2, m + m' \rangle, \end{aligned} \quad (40)$$

where

$$b_{-+} = \left(\frac{[2j][2]}{[2j + 1]} \right)^{1/2} q^{-1/2}, \quad (41)$$

$$b_{--} = - \left(\frac{[2j - 1]}{[2j + 1]} \right)^{1/2} q^{-j-1}. \quad (42)$$

As for the action of J_m , we get the following relations:

$$\begin{aligned} J_m |j, j + 1/2, m' \rangle = & a_{++} \begin{bmatrix} 1 & j + 1/2 & j + 1/2 \\ m & m' & m + m' \end{bmatrix}_q |j, j + 1/2, m + m' \rangle \\ & + a_{+-} \begin{bmatrix} 1 & j + 1/2 & j - 1/2 \\ m & m' & m + m' \end{bmatrix}_q |j, j - 1/2, m + m' \rangle. \end{aligned} \quad (43)$$

$$\begin{aligned}
J_m |j, j - 1/2, m' \rangle = & a_{-+} \begin{bmatrix} 1 & j - 1/2 & j + 1/2 \\ m & m' & m + m' \end{bmatrix}_q |j, j + 1/2, m + m' \rangle \\
& + a_{--} \begin{bmatrix} 1 & j - 1/2 & j - 1/2 \\ m & m' & m + m' \end{bmatrix}_q |j, j - 1/2, m + m' \rangle, \quad (44)
\end{aligned}$$

where the coefficients can be obtained as

$$a_{++} = \frac{[2j][2j+3]^{1/2}}{[2][2j+1]^{1/2}}, \quad a_{+-} = \frac{[2j+2]^{1/2}}{[2j+1]^{1/2}[2]^{1/2}}, \quad (45)$$

$$a_{-+} = -\frac{[2j]^{1/2}}{[2j+1]^{1/2}[2]^{1/2}}, \quad a_{--} = \frac{[2j+2][2j-1]^{1/2}}{[2][2j+1]^{1/2}}. \quad (46)$$

From these relations, we can define the operator representing total angular momentum as an operator which does not change the total spin. Indeed by defining the operator J_m^{tot} as

$$J_m^{tot} = q^{-1/2} J_m + \frac{1}{[2]} \sigma_m, \quad (47)$$

and using (37)(40)(43) and (44), we obtain

$$\begin{aligned}
J_m^{tot} |j, j + 1/2, m' \rangle &= \\
\left(\frac{[2j+1][2j+3]}{[2]^2} \right)^{1/2} \begin{bmatrix} 1 & j + 1/2 & j + 1/2 \\ m & m' & m + m' \end{bmatrix}_q |j, j + 1/2, m + m' \rangle, \quad (48)
\end{aligned}$$

$$\begin{aligned}
J_m^{tot} |j, j - 1/2, m' \rangle &= \\
\left(\frac{[2j-1][2j+1]}{[2]^2} \right)^{1/2} \begin{bmatrix} 1 & j - 1/2 & j - 1/2 \\ m & m' & m + m' \end{bmatrix}_q |j, j - 1/2, m + m' \rangle. \quad (49)
\end{aligned}$$

From these relations, we can find the value of the second order Casimir operator $C^{tot} = J^{tot} \cdot J^{tot}$ as

$$C^{tot} |j, j + 1/2, m \rangle = \frac{[2j+1][2j+3]}{[2]^2} |j, j + 1/2, m \rangle, \quad (50)$$

$$C^{tot} |j, j - 1/2, m \rangle = \frac{[2j-1][2j+1]}{[2]^2} |j, j - 1/2, m \rangle, \quad (51)$$

Note also that the operator O defined in (30) is no longer an identity operator. We obtain

$$O |j, j + 1/2, m \rangle = q^j |j, j + 1/2, m \rangle, \quad (52)$$

$$O |j, j - 1/2, m \rangle = q^{-j-1} |j, j - 1/2, m \rangle, \quad (53)$$

whereas $O'|j, j \pm 1/2, m \rangle = |j, j \pm 1/2, m \rangle$. We can also find that the action of the operator σ'_m on states can be written as $\sigma'_m = O^{-1}\sigma_m$.

Another Casimir operator can be constructed from the product of J_m and σ_m as $J \cdot \sigma$, which is the first-order operator with respect to the derivatives of the coordinates. The value of this Casimir operator can be obtained using relations (37), (40), (43) and (44):

$$\begin{aligned} J \cdot \sigma |j, j + 1/2, m \rangle &= (a_{++}b_{++} - b_{+-}a_{-+} + \frac{[2j]^{1/2}}{[2j+2]^{1/2}})|j, j + 1/2, m \rangle, \\ &= \frac{[2j]}{[2]}q^{j+1}, \end{aligned} \quad (54)$$

$$\begin{aligned} J \cdot \sigma |j, j - 1/2, m \rangle &= (a_{--}b_{--} - b_{-+}a_{+-} - \frac{[2j+2]^{1/2}}{[2j]^{1/2}})|j, j - 1/2, m \rangle \\ &= -\frac{[2j+2]}{[2]}q^{-j}|j, j - 1/2, m \rangle. \end{aligned} \quad (55)$$

Before trying to construct a Dirac operator, we first consider $\gamma_5(\gamma_3)$. In two dimensions, we usually take σ^i ($i = 1, 2$) as gamma matrices for the coordinates and σ_3 can be used as an γ_5 element. In our construction, we have already used all the Pauli matrices. This fact implies that one of the matrices we have used are redundant. As our γ_5 , we are going to choose the γ matrix corresponding to the direction of the radius. In other words, we take

$$\gamma_5 = iq^{-1} \frac{e \cdot \sigma'}{\sqrt{r^2}}, \quad (56)$$

where σ_m represents the Pauli matrix constructed from b'_m and c_m , and the normalization is chosen to ensure $\gamma_5^2 = -1$. This operator change the orbital angular momentum by ± 1 but does not change the total spin of the states. We therefore find

$$\gamma_5 |j, j + 1/2, m \rangle = \alpha_j |j + 1, j + 1/2, m \rangle, \quad (57)$$

$$\gamma_5 |j, j - 1/2, m \rangle = \beta_j |j - 1, j + 1/2, m \rangle, \quad (58)$$

where α_j and β_j are some coefficients which we do not need to specify for the foregoing discussion. Note also that this operator commutes with the total second order Casimir

operator C_{tot} . We are now going to construct Dirac operator. Our ansatz for the massless dirac operator is of the form

$$r\gamma \cdot \nabla = O^{-1}(J \cdot \sigma + a), \quad (59)$$

where r is the radius of the sphere. We will fix the value of a by requiring that this operator anticommute with γ_5 for states $|j, j + 1/2, m\rangle$ and $|j, j - 1/2, m\rangle$. It is not a priori clear that we can find the same a for these two states but we have $a = 1$. The value of the Dirac operator on the states is obtained as

$$r\gamma \cdot \nabla |j, j + 1/2, m\rangle = \frac{[2j + 2]}{[2]} |j, j + 1/2, m\rangle, \quad (60)$$

$$r\gamma \cdot \nabla |j, j - 1/2, m\rangle = -\frac{[2j]}{[2]} |j, j - 1/2, m\rangle. \quad (61)$$

From these we can find the relation to the total second order Casimir operator as

$$C^{tot} + \frac{1}{[2]^2} = (r\gamma \cdot \nabla)^2. \quad (62)$$

Note also that the massive Dirac operator $\gamma \cdot \nabla + im$ no longer anticommute with γ_5 .

We have constructed Dirac operator on the quantum two spheres. We expect that the fields with higher spins can be described using our approach. For physical application, the quantum four sphere is of particular interest. Indeed, there is no difficulty to extend the result of this paper to the quantum four spheres. We will discuss this issue in a separate paper[15].

References

- [1] V. G. Drinfeld, Sov. Math. Dokl. **32** (1985) 1254.
- [2] M. Jimbo, Lett. Math. Phys. **10** (1985) 63.
- [3] S. L. Woronowicz, Publ. RIMS, Kyoto Univ. **23** (1987) 117.
- [4] Yu. I. Manin, *Quantum Groups and Non-Commutative Geometry*, Preprint Montreal University, CRM-1561 (1988)
- [5] P. Podleś, Lett. Math. Phys. **14** (1987) 193
- [6] J. Wess, B.Zumino, Nuclear Physics B (Proc. Suppl.) **18B** (1990) 302.
- [7] U. Carow-Watamura, M. Schlieker, M. Scholl, S. Watamura, Z. Phys. C-Particles and Fields **48** (1990) 159; U. Carow-Watamura, M. Schlieker, S. Watamura, Comm. Math. Phys. **142** (1991) 605.
- [8] H.Suzuki, Prog. of Theor. Phys.91(1994)379.
- [9] N. Yu. Reshetikhin, L. A. Takhtadzhyan, and L. D. Faddeev, Leningrad Math. J. **1** (1990) 193.
- [10] A. N. Kirillov, N. Yu. Reshetikhin, *Representations of the Algebra $U_q(sl(2))$, q – Orthogonal Polynomials and Invariants of Links*, preprint 1988.
- [11] M.Noumi and K.Mimachi, Commun.Math.Phys.128(1990)521.
- [12] O.Ogievetsky, Lett.Math.Phys.24(1992),245.
- [13] O.Ogievetsky and B.Zumino, Lett.Math.Phys.25(1992),121.
- [14] G. Fiore, preprint SISSA-90-93-EP, hep-th/9403033
- [15] K.Ohta and H.Suzuki, in preparation.